

AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions and listings of claims in the application:

LISTING OF CLAIMS:

1. (currently amended): A fat emulsion with which a local anaesthetic is mixed before use, and which comprises propofol, an oily component, and an emulsifier, the fat emulsion further comprising a stabilizer selected from the following (a), (b), (c), or (d):

(a) at least one phospholipid selected from the group consisting of phosphatidylglycerol, phosphatidic acid, ~~phosphatidylinositol~~, and phosphatidylserine wherein a fatty acid esterified to a glycerol moiety is a C₁₈₋₂₂ linear or branched, saturated or unsaturated fatty acid;

(b) at least one phospholipid derivative selected from phosphatidylethanolamines modified with polyalkyleneglycol, wherein a fatty acid esterified to a glycerol moiety is a C₁₀₋₂₂ linear or branched, saturated or unsaturated fatty acid;

(c) at least one fatty acid selected from the group consisting of C₁₀₋₂₂ linear or branched, saturated or unsaturated fatty acids; or

(d) a mixture of at least two members selected from the above groups (a), (b), and (c), wherein

the stabilizer (a), (b), or (c) is present at a concentration of 0.01 to 1 w/v%, 0.01 to 1 w/v%, 0.05 to 5 w/v%, respectively, per the total amount of the fat emulsion and the local anaesthetic to be mixed therewith before use.

2. (previously presented): The fat emulsion according to Claim 1, wherein

(1) propofol is present at a concentration of 0.4 to 5 w/v%,

(2) the oily component is present at a concentration of 2 to 20 w/v%, and

(3) the emulsifier is present at a concentration of 0.4 to 5 w/v%, per the total amount of the fat emulsion and the local anaesthetic to be mixed therewith before use.

3. (previously presented): The fat emulsion according to Claim 1, wherein the local anaesthetic is at least one member selected from the group consisting of lidocaine, mepivacaine, bupivacaine, ropivacaine, dibucaine, procaine, procaine chloride, tetracaine and pharmacologically acceptable acid addition salts thereof.

4. (previously presented): The fat emulsion according to Claim 1, wherein the local anaesthetic is present at a concentration of 0.01 to 1 w/v%, per the total amount of the fat emulsion and the local anaesthetic to be mixed therewith before use.

5. (currently amended): The fat emulsion according to Claim 1, wherein the stabilizer is at least one phospholipid (a) selected from the group consisting of phosphatidylglycerol, phosphatidic acid, ~~phosphatidylinositol~~, and phosphatidylserine, wherein a fatty acid esterified to a glycerol moiety is a C₁₈₋₂₂ linear or branched, saturated or unsaturated fatty acid.

6. (canceled).

7. (currently amended): The fat emulsion according to Claim 1, wherein the stabilizer is at least one phospholipid (a) selected from the group consisting of distearoylphosphatidylglycerol, dioleoylphosphatidylglycerol, distearoylphosphatidic acid,

dioleoylphosphatidic acid, ~~distearoylphosphatidylinositol, dioleoylphosphatidylinositol,~~
distearoylphosphatidylserine, and dioleoyl-phosphatidylserine.

8. (previously presented): The fat emulsion according to Claim 5, wherein the stabilizer is present at a concentration of 0.03 to 1 w/v%, per the total amount of the fat emulsion and the local anaesthetic to be mixed therewith before use.

9. (original): The fat emulsion according to Claim 1, wherein the stabilizer is at least one phospholipid derivative (b) selected from phosphatidylethanolamines modified with polyalkyleneglycol, wherein a fatty acid esterified to a glycerol moiety is a C₁₀₋₂₂ linear or branched, saturated or unsaturated fatty acid.

10. (canceled).

11. (original): The fat emulsion according to Claim 1, wherein the stabilizer is at least one phospholipid derivative (b) selected from the group consisting of distearoylphosphatidylethanolamine-polyethylene glycol 5000, distearoylphosphatidylethanolamine-polyethylene glycol 3000, and distearoylphosphatidylethanolamine-polyethylene glycol 2000.

12. (previously presented): The fat emulsion according to Claim 9, wherein the stabilizer is present at a concentration of 0.1 to 1 w/v%, per the total amount of the fat emulsion and the local anaesthetic to be mixed therewith before use.

13. (original): The fat emulsion according to Claim 1, wherein the stabilizer is at least one fatty acid (c) selected from the group consisting of C₁₀₋₂₂ linear or branched, saturated or unsaturated fatty acids.

14. (canceled).

15. (original): The fat emulsion according to Claim 1, wherein the stabilizer is at least one fatty acid (c) selected from the group consisting of decanoic acid, lauric acid, myristic acid, palmitic acid, stearic acid, arachidic acid, isomyristic acid, isopalmitic acid, and oleic acid.

16. (previously presented): The fat emulsion according to Claim 13, wherein the stabilizer is present at a concentration of 0.1 to 5 w/v%, per the total amount of the fat emulsion and the local anaesthetic to be mixed therewith before use.

17. (previously presented): A fat emulsion containing container having a multi-compartment that is divided with a partition in such a manner as to allow contents in the compartments to come into contact with one another by opening the partition, which container comprising one compartment containing the fat emulsion according to Claim 1 and another compartment containing a local anaesthetic.

18. (currently amended): A pain-relieving fat emulsion comprising propofol, an oily component, an emulsifier, a stabilizer, and a local anaesthetic, wherein the stabilizer is selected from the following (a), (b), (c), or (d):

(a) at least one phospholipid selected from the group consisting of phosphatidylglycerol, phosphatidic acid, ~~phosphatidylinositol~~, and phosphatidylserine wherein a fatty acid esterified to a glycerol moiety is a C₁₈₋₂₂ linear or branched, saturated or unsaturated fatty acid;

(b) at least one phospholipid derivative selected from phosphatidylethanolamines modified with polyalkyleneglycol, wherein a fatty acid esterified to a glycerol moiety is a C₁₀₋₂₂ linear or branched, saturated or unsaturated fatty acid;

(c) at least one fatty acid selected from the group consisting of C₁₀₋₂₂ linear or branched, saturated or unsaturated fatty acids; or

(d) a mixture of at least two members selected from the above groups (a), (b), and (c), wherein

the stabilizer (a), (b), or (c) is present at a concentration of 0.01 to 1 w/v%, 0.01 to 1 w/v%, and 0.05 to 5 w/v%, respectively, in the fat emulsion.

19. (previously presented): The pain-relieving fat emulsion according to Claim 18, wherein

- (1) propofol is present at a concentration of 0.4 to 5 w/v%,
- (2) the oily component is present at a concentration of 2 to 20 w/v%,
- (3) the emulsifier is present at a concentration of 0.4 to 5 w/v%, and
- (4) the local anaesthetic is present at a concentration of 0.01 to 1 w/v%, in the fat emulsion.

20. (previously presented): The pain-relieving fat emulsion according to Claim 18, wherein the local anaesthetic is at least one member selected from the group consisting of

lidocaine, mepivacaine, bupivacaine, ropivacaine, dibucaine, procaine, procaine chloride, tetracaine and pharmacologically acceptable acid addition salts thereof.

21. (currently amended): The pain-relieving fat emulsion according to Claim 18, wherein the stabilizer is at least one phospholipid (a) selected from the group consisting of phosphatidylglycerol, phosphatidic acid, ~~phosphatidylinositol~~, and phosphatidylserine, wherein a fatty acid esterified to a glycerol moiety is a C₁₈₋₂₂ linear or branched, saturated or unsaturated fatty acid.

22. (canceled).

23. (currently amended): The pain-relieving fat emulsion according to Claim 18, wherein the stabilizer is at least one phospholipid (a) selected from the group consisting of distearoylphosphatidylglycerol, dioleoylphosphatidylglycerol, distearoylphosphatidic acid, dioleoylphosphatidic acid, ~~distearoylphosphatidylinositol~~, ~~dioleoylphosphatidylinositol~~, distearoylphosphatidylserine, and dioleoylphosphatidylserine.

24. (original): The pain-relieving fat emulsion according to Claim 21, wherein the stabilizer is present at a concentration of 0.03 to 1 w/v% in the fat emulsion.

25. (original): The pain-relieving fat emulsion according to Claim 18, wherein the stabilizer is at least one phospholipid derivative (b) selected from phosphatidylethanolamines modified with polyalkyleneglycol, wherein a fatty acid esterified to a glycerol moiety is a C₁₀₋₂₂ linear or branched, saturated or unsaturated fatty acid.

26. (canceled).

27. (original): The pain-relieving fat emulsion according to Claim 18, wherein the stabilizer is at least one phospholipid derivative (b) selected from the group consisting of distearoylphosphatidylethanolamine-polyethylene glycol 5000, distearoylphosphatidylethanolamine-polyethylene glycol 3000, and distearoylphosphatidylethanolamine-polyethylene glycol 2000.

28. (original): The pain-relieving fat emulsion according to Claim 25, wherein the stabilizer is present at a concentration of 0.1 to 1 w/v% in the fat emulsion.

29. (original): The pain-relieving fat emulsion according to Claim 18, wherein the stabilizer is at least one fatty acid (c) selected from the group consisting of C₁₀₋₂₂ linear or branched, saturated or unsaturated fatty acids.

30. (canceled).

31. (original): The pain-relieving fat emulsion according to Claim 18, wherein the stabilizer is at least one fatty acid (c) selected from the group consisting of oleic acid, isomyristic acid, isopalmitic acid, decanoic acid, lauric acid, myristic acid, palmitic acid, stearic acid, and arachidic acid.

32. (original): The pain-relieving fat emulsion according to Claim 29, wherein the stabilizer is present at a concentration of 0.05 to 0.2 w/v% in the fat emulsion.

33. (currently amended): A method for manufacturing a pain-relieving fat emulsion, the method comprising:

mixing a local anaesthetic with a fat emulsion comprising propofol, an oily component, an emulsifier, and a stabilizer, wherein the stabilizer is selected from the following (a), (b), (c), or (d):

(a) at least one phospholipid selected from the group consisting of phosphatidylglycerol, phosphatidic acid, ~~phosphatidylinositol~~, and phosphatidylserine wherein a fatty acid esterified to a glycerol moiety is a C₁₈₋₂₂ linear or branched, saturated or unsaturated fatty acid;

(b) at least one phospholipid derivative selected from phosphatidylethanolamines modified with polyalkyleneglycol, wherein a fatty acid esterified to a glycerol moiety is a C₁₀₋₂₂ linear or branched, saturated or unsaturated fatty acid;

(c) at least one fatty acid selected from the group consisting of C₁₀₋₂₂ linear or branched, saturated or unsaturated fatty acids; or

(d) a mixture of at least two members selected from the above groups (a), (b), and (c), to thereby obtain a fat emulsion, wherein

the stabilizer (a), (b), or (c) is present at a concentration of 0.01 to 1 w/v%, 0.01 to 1 w/v%, and 0.05 to 5 w/v%, respectively, in the fat emulsion.

34. (canceled).